

Anti-Rabbit IgG kappa, AlpSdAbs® VHH(iFluor488 ×4)

Summary

Code	025-104-007
Immunogen	Recombinant Rabbit IgG
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c
Conjugate	iFluor488(Ex: 495nm, Em: 519nm)
Specificity	Rabbit IgG kappa chain
Cross-Reactivity	No cross-reactivity with mouse, human, cynomolgus, rat, goat IgG
Purity	Recombinant Expression and Affinity purified
Concentration	1mg/ml
Formation	Liquid, 10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300, 50% glycerol
Storage	Store at -20 °C(Avoid freeze / thaw cycles) , Protect from light

Description

Anti-Rabbit IgG kappa, AlpSdAbs® VHH(iFluor488 ×4) is designed for detecting rabbit IgG kappa chain specifically. Anti-Rabbit IgG kappa, AlpSdAbs® VHH(iFluor488 ×4) is based on monovalent, recombinant single domain antibody to rabbit IgG kappa chain coupled to iFluor488. Based on immunoelectrophoresis and/or ELISA, Anti-Rabbit IgG kappa, AlpSdAbs® VHH(iFluor488 ×4) reacts with rabbit IgG kappa chain selectively, no reactivity with mouse, human, cynomolgus, rat, goat IgG.

Background

Rabbit research antibodies are widely used in life science research. So far, four isotypes have been identified (IgA, IgE, IgG, and IgM) in rabbits. Each isotype has a different heavy chain. Rabbit has only one IgG subclass. The whole IgG molecule possesses both the Fc region and the Fab region, which possessing the epitope-recognition site. The IgG contains two heavy and light chains. The heavy chain is about 50 KD and the light chain is about 25 KD. The common IgG is monomeric with a molecular weight of approximately 150 kD.

VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15KDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH. Thus VHH is considered of great value for research, diagnostics and therapeutics.

Benefits

High lot-to-lot consistency
 Increased sensitivity and higher affinity
 Animal-free production

Application notes

Flow Cyt	1:200-1:2000
ICC/IF	1:200-1:2000
ELISA	1:5000-1:20000
WB	1:5000-1:20000

Super-resolution microscopy

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical