

Anti-Rat IgG(H+L), AlpSdAbs® VHH(iFluor647 ×8)

Summary

Code	054-102-009
Immunogen	Rat IgG
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c
Conjugate	iFluor647(Ex:651nm, Em:667nm)
Specificity	Rat IgG(H+L)
Cross-Reactivity	Does not bind to mouse IgG, rabbit IgG, goat IgG, human IgG
Purity	Recombinant Expression and Affinity purified
Concentration	1mg/ml
Formation	Liquid, 10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300, 50% glycerol
Storage	Store at -20 °C(Avoid freeze / thaw cycles), Protect from light

Description

Anti-Rat IgG(H+L), AlpSdAbs® VHH(iFluor647 ×8) is designed for detecting rat IgG specifically. Anti-Rat IgG(H+L), AlpSdAbs® VHH(iFluor647 ×8) is based on recombinant single domain antibody to rat IgG coupled to iFluor647. Based on immunoelectrophoresis and/or ELISA, Anti-Rat IgG(H+L), AlpSdAbs® VHH(iFluor647 ×8) reacts with rat IgG(H+L) selectively, no reactivity with mouse IgG, rabbit IgG, goat IgG, human IgG.

Background

There are five antibody isotypes (IgA, IgD, IgE, IgG, and IgM) from rat. Each isotype has a different heavy chain. Rat IgG consists of four subclasses-IgG1, IgG2a, IgG2b, IgG2c. The whole IgG molecule possesses both the Fc region and the Fab region, which possessing the epitope-recognition site. The IgG contains two heavy and light chains, and the heavy chain is about 50 KD and the light chain is about 25 KD. The common IgG is monomeric with a molecular weight of approximately 150 kD.

VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15KDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH. Thus VHH is considered of great value for research, diagnostics and therapeutics.

Benefits

High lot-to-lot consistency
 Increased sensitivity and higher affinity
 Animal-free production

Application notes

Flow Cyt	1:200-1:2000
ICC/IF	1:200-1:2000
ELISA	1:5000-1:20000
WB	1:5000-1:20000
Super-resolution microscopy	

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical