

Anti-Mouse IgG2b(Fcy Fragment specific), AlpSdAbs[®] VHH(PE)

Code	001-104-010
Immunogen	Recombinant Fc region of mouse IgG2b
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c
Conjugate	PE(Ex: 488nm, Em: 575nm)
Specificity	Mouse IgG2b(Fcy fragment specific)
Cross-Reactivity	No cross-reactivity with mouse IgG1/2a/3, mouse IgM, rabbit, human, cynomolgus, rat, goat IgG
Purity	Recombinant Expression and Affinity purified
Concentration	0.1mg/mL
Formation	Liquid, 10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300
Storage	Store at 2-8 °C, Protect from light.

Description

Anti-Mouse IgG2b(Fcy Fragment specific), AlpSdAbs[®] VHH(PE) is designed for detecting mouse IgG2b Fcy fragment specifically. Anti-Mouse IgG2b(Fcy Fragment specific), AlpSdAbs[®] VHH(PE) is based on recombinant single domain antibody to mouse IgG2b Fc fused to PE. Based on immunoelectrophoresis and/or ELISA, Anti-Mouse IgG2b(Fcy Fragment specific), AlpSdAbs[®] VHH(PE) reacts with the Fc fragment of mouse IgG2b selectively, no reactivity with other mouse IgG subclasses, mouse IgM, or the Fab portion of mouse immunoglobulins.

Background

VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15KDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH.

Benefits

High lot-to-lot consistency
Increased sensitivity and higher affinity
Animal-free production

Application notes

Flow Cyt 1:200-1:1000
ICC/IF 1:200-1:1000

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

Please note: All products are FOR RESEARCH USE ONLY, NOT FOR USE IN DIAGNOSTIC PROCEDURES.